arXiv:1601.02691v1 [math.CA] 11 Jan 2016

An observation on "Classification of Lie point symmetries for quadratic Liénard type equation $\ddot{x}+f(x)\dot{x}^2+g(x)=0$" [J. Math. Phys. 54, 053506 (2013)] and its erratum [J. Math. Phys. 55, 059901 (2014)]

A Paliathanasis*[1] and PGL Leach†[2,3,4]

[1]Instituto de Ciencias Físicas y Matemáticas, Universidad Austral de Chile, Valdivia, Chile
[2]Department of Mathematics and Institute of Systems Science, Durban University of Technology, PO Box 1334, Durban 4000, Republic of South Africa
[3]School of Mathematics, Statistics and Computer Science, University of KwaZulu-Natal, Private Bag X54001, Durban 4000, Republic of South Africa
[4]Department of Mathematics and Statistics, University of Cyprus, Lefkosia 1678, Cyprus

January 13, 2016

Abstract

We demonstrate a simplification of some recent works on the classification of the Lie symmetries for a quadratic equation of Liénard type. We observe that the problem could have been resolved more simply.

In [1] and [2] the classification of the Lie (point) symmetries for the quadratic equation of the form

$$\ddot{x}+f(x)\dot{x}^2+g(x)=0, \tag{1}$$

was performed, where overdot denotes total differentiation with respect to time, "t".

We observe that under the coordinate (not point) transformation,

$$y=\int\left[\exp\left(\int f(x)\,dx\right)\right]dx, \tag{2}$$

equation (1) takes the simpler form

$$\ddot{y}+F(y)=0, \tag{3}$$

*anpaliat@phys.uoa.gr
†leach@ucy.ac.cy

where

$$g(x) = e^{-\int f(x)dx} F\left(\int e^{\int f(x)dx} dx\right). \tag{4}$$

The classification of the Lie symmetries of the latter equation has been known for some time (approximately 140 years). The various possibilities are

1. $F(y)$ is an arbitrary function. Equation (3) admits the autonomous symmetry, ∂_t, and the equation can be simply reduced to a first integral which in general cannot be evaluated to obtain the solutions in closed form.

2. In the two cases

   (a) $F(y) = (\alpha + \beta y)^n$, $n \neq 0, 1, -3$ and

   (b) $F(y) = e^{\gamma y}$, $\gamma \neq 0$,

   the admitted Lie symmetries of (3) constitute the algebra A_2 in the Mubarakzyanov Classification Scheme [3, 4, 5, 6].

3. When

   (a) $F(y) = \frac{1}{(y+c)^3}$ or

   (b) $F(y) = \alpha(y+c) + \frac{\beta}{(y+c)^3}$, $\beta \neq 0$,

   equation (3) is invariant under the three-dimensional algebra, $A_{3,8}$, which is more commonly known as $sl(2, R)$.

4. Finally for the cases

   (a) $F(y) = 0$,

   (b) $F(y) = c$,

   (c) $F(y) = y$ and

   (d) $F(y) = y + c$

   (note that in (c) and (d) a multiplicative constant – or arbitrary function of time which is beyond the considerations of [1, 2] – in the y term is superfluous), the algebra of the Lie symmetries is $sl(3, R)$ and, as a second-order linear equation, (3) is maximally symmetric [7][p 405].

Acknowledgements

AP acknowledges Prof. PGL Leach, Sivie Govinder, as also DUT for the hospitality provided and the UKZN, South Africa, for financial support. The research of AP was supported by FONDECYT postdoctoral grant no. 3160121.

References

[1] A.K. Tiwari, S.N. Pandey, M. Senthilvelan, and M. Lakshmanan, *Journal of Matheatical Physics* **54**, 053506 (2013)

[2] A.K. Tiwari, S.N. Pandey, M. Senthilvelan, and M. Lakshmanan, *Journal of Matheatical Physics* **55**, 059901 (2014)

[3] V.V. Morozov, Classification of six-dimensional nilpotent Lie algebras, *Izvestia Vysshikh Uchebn Zavendeniĭ Matematika,* **5** 161 (1958)

[4] G.M., Mubarakzyanov, On solvable Lie algebras, *Izvestia Vysshikh Uchebn Zavendeniĭ Matematika,* **32** 114 (1963)

[5] G.M. Mubarakzyanov, Classification of real structures of five-dimensional Lie algebras, *Izvestia Vysshikh Uchebn Zavendeniĭ Matematika,* **34** 99 (1963)

[6] G.M. Mubarakzyanov, Classification of solvable six-dimensional Lie algebras with one nilpotent base element, *Izvestia Vysshikh Uchebn Zavendeniĭ Matematika,* **35** 104 (1963)

[7] S.M. Lie *Differentialgleichungen* (Chelsea, New York, 1967)